



Turf and Ornamental Herbicide

Turf Fertilizer with Dimension® 0.17% FG (19-0-5)/30% Uflexx

For season-long control of crabgrass and control or suppression of listed annual grasses and broadleaf weeds in established lawns and ornamental turf, including golf course fairways, roughs and tee boxes.

- Not for use on plants beings grown for sale or other commercial use, or for commercial seed production, or for research purposes.
• For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns.

This product is NOT for sale in the state of New York.

GUARANTEED FERTILIZER ANALYSIS

Table with 2 columns: Component and Percentage. Includes Total Nitrogen (N) 19%, Available Phosphate (PO) 0%, Soluble Potash (KO) 5%, Magnesium (Mg) 4.326%, Sulfur (S) 0.300%, Iron (Fe) 3.00%.

Active ingredients:

Table with 2 columns: Ingredient and Percentage. Includes Dithiopyr: 3, 5- pyridinedicarbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-6-(trifluoromethyl)-S, S-dimethyl ester 0.17%, Inert Ingredients 99.83%, Total 100.00%.

Source of nutrients derived from: Urea, Urea N-(n-Butyl) thiophosphoric Triam, Dicyandiamide, Muriate of Potash, Iron (II) Oxide, Iron Sulfate, Dolomitic Limestone

NET WEIGHT: 50 LBS 22.68 KGS

Keep Out of Reach of Children CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals CAUTION: Causes Eye Irritation • Harmful if inhaled

Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

First Aid

If in eyes: hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move the person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Hot Live Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

Environmental Hazards

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters

Dimension® is the registered trademark of Dow AgroSciences LLC 9330 Zionsville Road, Indianapolis, IN 46268

Produced and Guaranteed By: Bonus Crop Fertilizer, Inc. PO Box 1062 Greenville, Texas 75403-1062 EPA Est. # 49991-TX-2 EPA Reg # 62719-504-4991

Notice: Read the entire label. Use only according to label directions. Before using this product, read Limited Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of Directions for use. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. Agricultural Chemical Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuff, feed or seed by storage or disposal.

Pesticide Storage: Store this product only in its original container in a dry, cool, secured storage area.

Pesticide Disposal: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available, or dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Use Directions for Turf

BCF Turf Fertilizer with Dimension® 0.15% FG Turf herbicide provides season-long control of crabgrass and control or suppression of other listed annual grasses and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. When used according to label directions, this product will not harm nearby established ornamentals identified as "Tolerant Ornamentals". However, this product has not been tested on all ornamental species, all cultivars of species or under all possible growing conditions.

Use Precaution for Turf

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. Initial application of this product should be made only after turfgrasses have developed a vigorous root system, a uniform stand and have received at least two mowings following seeding, sodding or sprigging. Use of this product on turfgrasses that are not well-established, or weakened by stress from weather, pests, disease, chemical, or mechanical influences may increase the chances of turf injury.

Apply this product only to turfgrass species listed on this label. The following turfgrass species are tolerant to this product when applied as directed:

Cool-Season Grasses

Table with 2 columns: Grass Name and Scientific Name. Includes Bentgrass, creeping; Bluegrass, Kentucky; Fescue, fine; Fescue, tall; Ryegrass, perennial; Agrostis Palustris; Poa Pratensis; Festuca rubra; Festuca arundinacea; Lolium perenne.

Warm-Season Grasses

Table with 4 columns: Grass Name, Scientific Name, Grass Name, Scientific Name. Includes Bahiagrass; Bermudagrass; Buffalograss; Carpetgrass; Paspalum notatum; Cynodon Dactylon; Buchloe dactyloides; Axonopus affinis; Centipedegrass; Kikuyugrass; St. Augustinegrass; Zoysiagrass; Eremochloa ophiuroides; Pennisetum clandestinum; Stenotaphrum secundatum; Zoysia japonica.

Use of this product on certain varieties of creeping bentgrass, such as 'Cohansey', 'Carmen', 'Seaside', and 'Washington' may result in undesirable turfgrass injury. Not all varieties of creeping bentgrass have been tested. **Do not** apply this product to Colonial bentgrass (*Agrostis tenuis*).

Use of the product on certain varieties of fine fescue may result in undesirable turf injury. The following fine fescue varieties have been found to be sensitive to this product: 'Atlanta', 'Banner', 'Beauty', 'Bigart', 'CF-2', 'Enjoy', 'HF-93', 'Highlight', 'Ivalo', 'Jamestown', 'Koker', 'Majenta', 'Mary', 'Pennlawn', 'Tamara', 'Tajana', 'Waldorl', and 'Waldina.' Not all varieties of fine fescue have been tested.

Do not use this product on seeding Buffalograss in the spring of the first year of establishment until the turfgrass is fully green and has established new roots.

Note: It is not possible to evaluate all turfgrass varieties for tolerance to this product. It is suggested that the user evaluate the response of turfgrass varieties not listed on this label on a limited area, at recommended use rates, prior to initiating large-scale use.

Precautions for Reseeding, Overseeding, Sprigging

- Reseeding, overseeding, or sprigging of areas treated with this product within 10 weeks after a single application of this product or within 16 weeks after a split application program totaling 7.7lb/1000sq ft (0.5 lb active ingredient per acre) or more may inhibit the establishment of desirable turfgrasses.
- When reseeding or overseeding, follow established cultural practices for seedbed preparation, seeding, irrigation and fertilization.
- For best overseeding results, use mechanical or power seeding equipment (sill seeders) designed to provide good contact between soil and seed.

Precautions for controlling Crabgrass

Early postemergence weed control with this product is limited to crabgrass. Postemergence application effectively controls crabgrass, but only if application is made prior to the 3-leaf stage of growth or first tiller. For best results, cultural practices that disturb the soil, such as core-, spike-, or hydro-aerification, and verticulating, should be done before application of this product. **Do not** apply this product until turfgrass has recovered from the mechanical injury caused by these cultural practices.

Other Use Precautions

- **Do not** use clipping from treated turf for mulching around vegetables or fruit trees.
- **Do not** allow people or pets enter treated area until dust has settled.
- **Do not** graze livestock or feed forage cut from areas treated with this product.

Application Directions for Turf

Tips for Improved Weed Control

- For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application.
- The turf herbicide and fertilizer components in this product are not effective until activated by rainfall or irrigation. Performance is improved if application is followed by ½ inch of rainfall or sprinkler irrigation. Erratic weed control may result if not activated by rainfall or irrigation within 30 days after application.
- Use of split (half-rate) applications spaced 5-10 weeks apart may provide improved weed control.

Maximum Use Rates (Applies to all Turfgrass Uses)

- **Do not** apply more than 333lbs. (0.5lb active ingredient) per acre (7.7lb per 1000 sq ft) per application or more than 1000 lb (1.5 lb active ingredient) per acre (23.1lb per 1000 sq ft) per year if using split or sequential applications.

Control of Crabgrass

Preemergence and Early Postemergence Control

- This product provides "preemergence" control of crabgrass in established turfgrasses (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass.
- This product also provides "early postemergence" control of crabgrass if applied prior to the 3-leaf stage of growth or first tiller as described below. This stage of growth generally corresponds to the time when crabgrass seedlings become visible in the established turfgrass. The practical benefit of early postemergence crabgrass control is a 2-4 week longer period of time (window of application) during which crabgrass control can be achieved.

Application Frequency and Timing

This product may be applied as a single application, as a split application, or as a sequential application for crabgrass control in the spring, summer, or fall as described in the following use programs.

Spring Application

For application made in the spring or early summer, use an appropriate application rate corresponding to one of the three control programs listed in Table 1. Specific rate recommendations are based on the user's location (Region), the mowing height of the turfgrass and whether the application timing is considered to be preemergence or early postemergence. The duration of residual weed control provided by this product is directly related to application rate, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use Program 1 for preemergence control at sites where turfgrass is cut relatively high (e.g. homeowner lawns). This program provides 3-5 months of preemergence crabgrass control. This program will also provide early postemergence control of crabgrass up to the 3-leaf stage of first tiller at sites where turfgrass is cut relatively high (e.g. homeowner lawns)

Use Program 2 for preemergence control at sites where (a) turfgrass is cut relatively low (e.g. golf fairways) and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g. residential lawns).

Use Program 3 for preemergence control at sites where (a) turfgrass is cut relatively low (e.g. golf fairways) and (b) turfgrass maintenance or weed control has been conducted during the previous year. This program provides 4-6 months of preemergence crabgrass control. This program may also be used for early postemergence control of crabgrass up to the 3-leaf stage or first tiller at sites where turfgrass is cut relatively high (e.g. golf fairways).

Table 1 Recommended Use Rates

Region	Application Rates	Program 1	Program 2	Program 3
All States, except NY, and parts of states not include in transition, south, coastal south or west	Ib/acre	127	167	253
	Lb/1000 sq ft	2.9	3.8	5.8
	Lb ai/acre	0.19	0.25	0.38
Transition; DE, KS, KY, MD, MO, NJ ,VA, southeastern PA, southern areas of IL, IN, OH & coastal areas of ct, & RI	Ib/acre	167	253	333
	Lb/1000 sq ft	3.8	5.8	7.7
	Lb ai/acre	0.25	0.38	0.5
South; AL, AR, GA, LA, MS, NC, NM, OK, SC, TN, & TX	Ib/acre	253	167+167	207+207
	Lb/1000 sq ft	5.8	3.8+3.8	4.7+4.7
	Lb ai/acre	0.38	0.25+0.25	0.31+0.31
Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX	Ib/acre	167+167	207+207	253+253
	Lb/1000 sq ft	3.8+3.8	4.7+4.7	5.8+5.8
	Lb ai/acre	0.25+0.25	0.31+0.31	0.38+0.38
West: AZ, CA, & NV. In this climatically diverse region, use the higher rates in local areas with longer crabgrass seasons	Ib/acre	127+167	167+253	167+167
	Lb/1000 sq ft	2.9+3.8	3.8+5.8	3.8+3.8
	Lb ai/acre	0.19+0.25	0.25+0.38	0.25+0.25

Split applications may be made 5-10 weeks apart

Use of Split or Sequential Applications: Split or sequential applications may be used to provide improved weed control or to provide extended weed control in areas with long growing season. Split applications may be made with the rates in Table 1 being split between two applications made 5-10 weeks apart. Where longer periods of control are required, sequential applications may be made in late summer or fall following a spring application or in the spring following a fall application, provided the maximum application rate per year is not exceeded. Refer to the "Maximum Use Rates" section above.

Terms and Conditions of Use: If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the Seller for a full refund of purchase price paid.

Warranty Disclaimer: Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OR MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use: It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.) abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies: The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following: 1) Refund of purchase price paid by Buyer or User for product bought, or 2) Replacement of amount of product used. The terms of the Warranty Disclaimer, Inherent Risks of Use and the Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfc.org/metals.htm>

SAFETY DATA SHEET



BCF TURF FERTILIZER with DIMENSION

Emergency Phone: 800-992-5994
Bonus Crop Fertilizer
Greenville, TX 75403

Effective Date: 5/28/2015

1. PRODUCT AND COMPANY IDENTIFICATION:

PRODUCT: Dimension on Fertilizers

COMPANY IDENTIFICATION:

Bonus Crop Fertilizer, Inc.
P.O Box 1062
Greenville, TX 75403

2. COMPOSITION/INFORMATION ON INGREDIENTS:

Dithiopyr:	CAS # 097886-45-8	0.06- 0.25%
S,S'-dimethyl 2-(difluoromethyl-4-(2-methylpropyl)-6-(trifluoromethyl)-3,5-pyridinedicarbothioate		
Other Ingredients, Total, Including Silica		99.75-99.94%
Dipropylene glycol Fertilizers		

CAUTION



This document is prepared pursuant to the OSHA Hazard Communication Standard (29 CFR 1910.1200). In addition, other substances not 'Hazardous' per this OSHA Standard may be listed. Where proprietary ingredient shows, the identity may be made available as provided in this standard.

3. HAZARDOUS IDENTIFICATIONS:

EMERGENCY OVERVIEW

Color Appearance Odor. May cause eye irritation. Toxic to aquatic organisms.

EMERGENCY PHONE NUMBER: 800-992-5994

POTENTIAL HEALTH EFFECTS: This section includes possible adverse effects, which could occur if this material is not handled in the recommended manner.

EYE: May cause eye irritation.

SKIN: Prolonged exposure is not likely to cause significant skin irritation. Prolonged skin contact is unlikely to result in absorption of harmful amounts. The dermal LD₅₀ has not been determined. Dithiopyr did not cause allergic skin reactions when tested in guinea pigs.

INGESTION: Very low toxicity if swallowed. Harmful effects not anticipated from swallowing small amounts. Single dose oral LD₅₀ has not been determined. Based on information for the components, the oral LD₅₀ for rats is expected to be >5000 mg/kg.

INHALATION: At room temperature, exposure to vapor is minimal due to low volatility; single exposure is not likely to be hazardous.

SYSTEMIC (OTHER TARGET ORGAN) EFFECTS:

Dithiopyr, in animals, effects have been reported on the following organs: blood, kidney, liver, thyroid, adrenal gland and gall bladder.

CANCER INFORMATION: Dithiopyr did not cause cancer in laboratory animals.

TERATOLOGY: Dithiopyr did not cause birth defects or other effects in the fetus even at doses which caused toxic effects in the mother.

REPRODUCTIVE EFFECTS: In animal studies, dithiopyr has been shown not to interfere with reproduction.

4. FIRST AID:

EYE: Flush eyes thoroughly with water for several minutes. Remove contact lenses after initial 1-2 minutes and continue flushing for several additional minutes. If effects occur, consult a physician, preferably an ophthalmologist.

SKIN: Wash skin with plenty of water.

INGESTION: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Never give anything by mouth to an unconscious person.

INHALATION: Move person to fresh air; if effects occur, consult a physician.

NOTE TO PHYSICIAN: No specific antidote. Treatment of exposure should be directed at the control of symptoms and the clinical condition of the patient.

SAFETY DATA SHEET



BCF TURF FERTILIZER with DIMENSION

Emergency Phone: 800-992-5994
Bonus Crop Fertilizer
Greenville, TX 75403

Effective Date: 5/28/2015



maximum recommended storage temperature for this material is 95°F (35°C).

5. FIRE FIGHTING MEASURES:

FLASH POINT: 363°F (184°C)

FLAMMABLE LIMITS:

LFL: Not determined

ULF: Not determined

EXTINGUISHING MEDIA: Use the following extinguishing media when fighting fires involving this material; carbon dioxide, dry chemical or water spray

UNUSUAL HAZARDS: Pesticide particulates can become airborne.

FIRE-FIGHTING EQUIPMENT: Wear self-contained breathing apparatus (pressure demand NIOSH approved or equivalent) and full protective gear.

6. ACCIDENTAL RELEASE MEASURES:

ACTION TO TAKE FOR SPILLS/LEAKS: Sweep up small spills and place in a suitable container for disposal. Report large spills to 800-992-5994. Keep dust to a minimum. Keep out of sewers and open bodies of water.

Wear appropriate protective equipment when handling a spill of this material. If exposed to material during clean-up operations see Section 4 for actions to follow. Remove all contaminated clothing promptly. Wash all exposed skin areas with soap and water immediately after exposure. Thoroughly launder clothing before reuse. Do not take clothing home to be laundered.

7. HANDLING AND STORAGE:

PRECAUTIONS TO BE TAKEN IN HANDLING AND STORAGE: Keep out of reach of children. Avoid skin and eye contact. Wash thoroughly with soap and water after handling. Do not swallow. Do not store this material near food, feed, or drinking water. The

8. EXPOSURE CONTROLS/PERSONAL PROTECTION:

These precautions are suggested for conditions where the potential for exposure exists. Emergency conditions may require additional precautions.

EXPOSURE GUIDELINES: None established.

ENGINEERING CONTROLS: Good general ventilation should be sufficient for most conditions. Local exhaust ventilation may be necessary for some operations.

RECOMMENDATIONS FOR MANUFACTURING, COMMERCIAL BLENDING, AND PACKAGING WORKERS:

EYE/FACE PROTECTION: Use safety glasses.

SKIN PROTECTION: No precautions other than clean body-covering clothing should be needed.

RESPIRATORY PROTECTION: No respiratory protection should be needed.

APPLICATORS AND ALL OTHER HANDLERS: Refer to the product label for personal protective clothing and equipment.

9. PHYSICAL AND CHEMICAL PROPERTIES:

COLOR: Tan

STATE: Solid

ODOR: Sulfur

BOILING POINT: Not applicable

MELTING POINT: 149°F (65°C)

VAPOR PRESSURE: Negligible

DENSITY: 1.41 g/cm³

SOLUBILITY IN WATER: 1.38 ppm @ 68°F (20°C)

SPECIFIC GRAVITY: (Water = 1): Not determined

PERCENT VOLATILITY: 0%

pH: 4.2

VISCOSITY: Not applicable

EVAPORATION RATE: (Bac = 1): Not applicable

10. STABILITY AND REACTIVITY:

SAFETY DATA SHEET



BCF TURF FERTILIZER with DIMENSION

Emergency Phone: 800-992-5994
Bonus Crop Fertilizer
Greenville, TX 75403

Effective Date: 5/28/2015

STABILITY: This material is stable under normal storage conditions.

INCOMPATIBILITY: None known

HAZARDOUS DECOMPOSITION PRODUCTS: None known

HAZARDOUS POLYMERIZATION: Not known to occur.

11. TOXICOLOGICAL INFORMATION:

MUTAGENICITY: For the components tested, in-vitro and animal mutagenicity studies were negative.

12. ECOLOGICAL INFORMATION:

ENVIRONMENTAL FATE:

MOVEMENT & PARTITIONING:

Based largely or completely on information for the dipropylene glycol.

Bioconcentration potential is low (BCF is <100 or Log Pow <3).

Based largely or completely on information for the silica. Partitioning from water to n-octanol is not applicable.

DEGRADATION & PERSISTENCE:

Based largely or completely on information for the dipropylene glycol.

Based on stringent OECD test guidelines, this material cannot be considered as readily biodegradable; however, these results do not necessarily mean that the material is not biodegradable under environmental conditions.

Material is inherently biodegradable. Reaches more than 20% biodegradation in OECD test(s) for inherent biodegradability.

Based largely or completely on information for the silica. Biodegradation is not applicable.

ECOTOXICOLOGY:

Based largely or completely on information for the active ingredient.

Material is highly toxic to aquatic organisms on an acute basis (LC₅₀ or EC₅₀ is between 0.1 and 1 mg/L in the most sensitive species tested).

Material is practically non-toxic to birds on an acute basis (LD₅₀ is >2000 mg/kg).

Material is practically non-toxic to birds on a dietary basis (LC₅₀ is >5000 ppm).

Based largely or completely on information for the dipropylene glycol.

Material is practically non-toxic to aquatic organisms on an acute basis (LC₅₀ or EC₅₀ is >1000 mg/L in the most sensitive species tested).

Based largely or completely on information for the silica. Not expected to be acutely toxic to aquatic organisms.

13. DISPOSAL CONSIDERATIONS:

DISPOSAL METHOD: Wastes resulting from the use of this product may be disposed of on site according to label directions or at an approved waste disposal facility. Follow all local, state and federal requirements for disposal.

14. TRANSPORT INFORMATION:

U.S. DEPARTMENT OF TRANSPORTATION (DOT) INFORMATION:

For non-bulk shipments by land: Material is not regulated for transportation.

For bulk shipments by land or vessel: ENVIRONMENTALLY HAZARDOUS SUBSTANCES, SOLID, N.O.S. (DITHIOPYR)/9/UN3077PGIII/MARINE POLLUTANT (DITHIOPYR)

15. REGULATORY INFORMATION:

NOTICE: The information herein is presented in good faith and believed to be accurate as of the effective date shown above. However, no warranty, express or implied, is given. Regulatory requirements are subject to change and may differ from one location to another; it is the buyer's responsibility to ensure that its activities comply with federal, state or provincial, and local laws. The following specific information is made for the purpose of complying with numerous federal, state or provincial, and local laws and regulations.

U.S. REGULATIONS

SARA 313 INFORMATION: To the best of our knowledge, this product contains no chemical subject to SARA Title III Section 313 supplier notification requirements.

SARA HAZARD CATEGORY: This product has been reviewed according to the EPA "Hazard Categories"

SAFETY DATA SHEET



BCF TURF FERTILIZER with DIMENSION

Emergency Phone: 800-992-5994
Bonus Crop Fertilizer
Greenville, TX 75403

Effective Date: 5/28/2015

promulgated under Sections 311 and 312 of the Superfund Amendment and Reauthorization Act of 1986 (SARA Title III) and is considered, under applicable definitions, to meet the following categories:

An immediate health hazard
A delayed health hazard

TOXIC SUBSTANCES CONTROL ACT (TSCA): All ingredients are on the TSCA inventory or are not required to be listed on the TSCA inventory.

STATE RIGHT-TO-KNOW: This product is not known to contain any substances subject to the disclosure requirements of
New Jersey
Pennsylvania

OSHA HAZARD COMMUNICATION STANDARD: This product is a "Hazardous Chemical" as defined by the OSHA Hazard Communication Standard, 29 CFR 1910.1200.

COMPREHENSIVE ENVIRONMENTAL RESPONSE COMPENSATION AND LIABILITY ACT (CERCLA, or SUPERFUND): To the best of our knowledge, this product contains no chemical subject to reporting under CERCLA.

16. OTHER INFORMATION:

SDS STATUS: New 05/28/2015

The Information Herein Is Given In Good Faith, But No Warranty, Express or Implied, Is Made. Consult Bonus Crop Fertilizer @ 800-336-2434 for Further Information

Bonus Crop Fertilizer, Inc. urges each customer or recipient of this (M)SDS to study it carefully and consult appropriate expertise, as necessary or appropriate, to become aware of and understand the data contained in this (M)SDS and any hazards associated with the product. The information herein is provided in good faith and believed to be accurate as of the effective date shown above. However, no warranty, express or implied, is given. Regulatory requirements are subject to change and may differ between various locations. It is the Buyer's/User's responsibility to ensure that his activities comply with all federal, state, provincial or local laws. The information presented here pertains only to the product as shipped. Since conditions for use of the product are not under the control of the manufacturer, it is the Buyer's/User's duty to determine the conditions necessary for the safe use of this product. Due to the proliferation of sources for information such manufacturer-specific (M)SDS, we are not and cannot be responsible for (M)SDS obtained from any source other than ourselves. If you have obtained an (M)SDS from another source or if you are not sure that the (M)SDS you have is current, please contact us for the most current version.