

BroadStar™

HERBICIDE

GRANULAR PRE-EMERGENCE WEED CONTROL FOR NURSERY ORNAMENTALS



Successful weed control in nursery ornamental production begins with timely applications of the most effective herbicides. Once plants are established, a preemergence application of BroadStar™ Herbicide helps ensure clean containers and is an excellent resistance management tool to maximize the efficacy of your granular herbicide program. Plus, the proven performance of BroadStar provides the longest residual control of a broad spectrum of tough weeds.

BROADSTAR DELIVERS


- Proven granular preemergence weed control to avoid costly hand weeding
- Longest residual performance - 8 to 12+ weeks of control
- Broad spectrum control of tough broadleaf and grassy weeds including spurge, bittercress, eclipsta and winter annuals such as chickweed, groundsel, liverwort and many more

APPLICATION INFORMATION

APPLICATION RATE	APPLICATION TIMING	GRANULE ACTIVATION
150 lbs/A (approximately 3.5 lbs per 1,000 sq ft)	<ul style="list-style-type: none"> • Apply to established woody ornamentals at least 6 weeks after transplanting • Apply prior to weed emergence 	Apply 1/2 to 3/4 inch of irrigation by hand, overhead sprinkler or fan emitter after application to ensure activation and effective weed control

NOTE: For effective resistance management, be sure to alternate BroadStar with other preemergence chemistries.

BROADSTAR DELIVERS OUTSTANDING, LONG-LASTING WEED CONTROL



Dr. Jeff Derr, Virginia Tech – Treated on 11/7; Results on 12/2 – Irrigated 1/2" of water immediately after application



www.nufarm.com/us

©2015 Nufarm. Important: Always read and follow label instructions. BroadStar® is a trademark of Valent U.S.A. Corporation. OH2® is a trademark of OMS Investments, Inc. Regal 0-0® is a trademark of Regal Chemical Company. Ronstar® is a trademark of Bayer CropScience. Rout® is a trademark of Scotts-Sierra Crop Protection Company. 15-GHN-0022



Grow a better tomorrow.